

A Technique for Measuring Electrical Currents in the Solar Corona

Steven R. Spangler

Department of Physics and Astronomy, University of Iowa, Iowa City, IA 52242

ABSTRACT

A technique is described for measuring electrical currents in the solar corona. It uses radioastronomical polarization measurements of a spatially-extended radio source viewed through the corona. The observations yield the difference in the Faraday rotation measure between two closely-spaced lines of sight through the corona, a measurement referred to as *differential Faraday rotation*. It is shown that the expression for differential Faraday rotation is proportional to the path integral $\oint n \vec{B} \cdot d\vec{s}$ where n is the plasma density and $\vec{B}$ is the coronal magnetic field. The integral is around a closed loop (Amperian Loop) in the corona. If the plasma density is assumed roughly constant, the differential Faraday rotation is proportional to the current within the loop, via Ampere's Law. The validity of the constant density approximation is discussed, and two test cases are presented in which the associated error in the inferred current is small, of order tens of percent or less. The method is illustrated with observations of the radio source 3C228 with the Very Large Array (VLA) in August, 2003. A measurement of a differential Faraday rotation "event" on August 16, 2003, yields an estimate of 2.5×10^9 Amperes in the Amperian Loop. A smaller event on August 18 yields an enclosed current of 2.3×10^8 Amperes. The implications of these currents for coronal heating are briefly discussed.

Subject headings: Sun:corona—Sun:magnetic fields—plasmas

1. Introduction

Electrical currents certainly flow in the solar corona. The structure seen in eclipse photographs or coronagraph images shows that pressure gradients must be balanced by electrodynamic forces. Although information on the strength and form of the coronal magnetic field is limited, it is clear that the true field is deformed from the potential field generated by a magnetic scalar potential. This difference is due to the presence of electrical currents.

In addition, there is a class of theories for heating of the solar corona which invoke Joule heating from coronal currents, probably contained in turbulent current sheets. This idea originated with Parker (1972), and has been elaborated in many subsequent works (see, for example, Gudiksen and Nordlund 2005; Peter et al 2006).

Observational measurements of coronal currents appear to be nonexistent. In fact, measurements of the coronal magnetic field itself at heliocentric distances of a few solar radii are limited to results from Faraday rotation of a trans-coronal radio source and analysis of radio emission from solar flares (Bird and Edenhofer 1990). There is a literature on results from Faraday rotation observations of the large scale structure of the coronal field, as well as magnetic field inhomogeneities on a wide range of scales. Examples of such papers, which give references to the wider literature are Hollweg et al (1982); Bird and Edenhofer (1990); Mancuso and Spangler (1999, 2000); Spangler (2005) and Ingleby et al (2007).

In this paper, I discuss how Faraday Rotation measurements can also provide an observational estimate of electrical currents in the corona. The technique requires measurements of *differential Faraday Rotation* (Spangler 2005), which is the difference in the Faraday rotation measure between two closely-spaced lines of sight through the corona. As discussed in the previously cited papers, a Faraday Rotation measurement yields the rotation measure, RM , given by Kraus (1966)

$$RM = \left(\frac{e^3}{8\pi^2 c^3 \epsilon_0 m_e^2} \right) \int_{LOS} n_e \vec{B} \cdot d\vec{z} \quad (1)$$

The fundamental physical constants of e, m_e, c , and ϵ_0 are, respectively, the fundamental charge, the mass of an electron, the speed of light, and the permittivity of free space. The electron density in the plasma is n_e , and $\vec{B}$ is the vector magnetic field. The incremental vector $d\vec{z}$ is a spatial increment along the line of sight, which is the path on which the radio waves propagate. Positive dz is in the direction from the source to the observer. The subscript LOS on the integral indicates an integral along the line of sight. Equation (1) is in SI units, as opposed to cgs, which has been used in our previous papers. The SI system is used in this paper for convenience in discussing electrical currents. The units of the rotation measure are radians/m².

As will be discussed in Section 2, when the rotation measures on two (or more) closely-spaced lines of sight are compared, one has an estimate of the electrical current between the two lines of sight. Such multiple lines of sight are available when one images an extended radio source (such as a radio galaxy or quasar) which is occulted by the corona. As is discussed in some of the papers referenced above, such measurements are straightforward

with the Very Large Array radiotelescope of the National Radio Astronomy Observatory¹.

The present investigation was motivated by laboratory experiments using differential Faraday rotation measurements to determine internal currents in the Madison Symmetric Torus (MST) Reversed Field Pinch (RFP) at the University of Wisconsin (Prager 1999) by D.L. Brower and W.X. Ding (Brower et al 2002; Ding et al 2003). In the case of the MST device, Ding and Brower use an infrared laser as the source of polarized radiation for measurement of Faraday Rotation. Measurement of the rotation measure along spatially-separated paths is achieved by directing the beam along distinct chords through the machine. In spite of obvious technical differences, the physical situation is virtually identical to that of the present paper. The inferred current profiles in the MST, and their dependence on time, are in agreement with theoretical predictions and the results from other diagnostics of the plasma. The results of Brower et al (2002) and Ding et al (2003) encourage application of this technique to the solar corona.

The organization of this paper is as follows. Section 2 describes the basis of the technique, i.e. how differential Faraday Rotation measurements can provide a measurement of electrical currents in the corona. Section 3 provides an observational implementation of this technique with Very Large Array observations of the radio source 3C228 in August, 2003 (Spangler 2005). In that section, it is shown that the total current contained within the Amperian Loop formed by the lines of sight was as high as 2.5 GigaAmperes during one period on August 16, but less than 0.8 GigaAmperes during a 3 hour period of high quality data before this event. A smaller, marginal detection of differential Faraday Rotation on August 18 yielded a current of the order of 0.23 GigaAmperes. In Section 4 we discuss the validity of an approximation used in the derivation of our expression for the current contained between two lines of sight through the corona. This assumption (stated in eq.[3] below) is that the plasma density can be approximated as roughly constant in the region that makes the maximum contribution to the Faraday rotation, and taken outside the integral defining the differential rotation measure. Arguments based on a simple theoretical model for coronal current sheets as well as data from a laboratory fusion plasma indicate that the error introduced by this approximation is relatively small, of order tens of percent or less. Section 5 briefly discusses the implications of our observations and analysis for theories of coronal heating by Joule dissipation of the currents. Our estimate of Joule heating using the Spitzer resistivity is about 6 orders of magnitude less than the observational estimate for the volumetric heating rate in the relevant part of the corona. Finally, Section 6 summarizes

¹The Very Large Array is an instrument of the National Radio Astronomy Observatory. The NRAO is a facility of the National Science Foundation, operated under cooperative agreement with Associated Universities, Inc.

and concludes.

2. Physical Basis of the Technique

The technique is based on simultaneous Faraday rotation measurements along two lines of sight through the corona, which are separated by a small angular distance on the sky, and a corresponding physical separation l in the solar corona. Such observations can and have been made with the Very Large Array (VLA) of radio galaxies and quasars (Sakurai and Spangler 1994; Mancuso and Spangler 1999; Spangler 2005). A specific illustration is our observation of the radio galaxy 3C228 on August 16 and August 18, 2003 (Spangler 2005; Spangler et al 2007). As may be seen in Figure 2 of Spangler (2005), 3C228 is a double radio source with bright, highly polarized hot spots separated by about 46 arcseconds on the sky. The corresponding physical separation between the lines of sight to the two hot spots in the corona is about 33,000 km. The observations of August 16 and August 18 were made when the lines of sight passed within $6.7R_{\odot}$ and $5.2R_{\odot}$ of the center of the Sun, respectively.

A Faraday Rotation measurement of an extended radio source with two components is illustrated in cartoon form in Figure 1(a). The radio telescope measures rotation measure values RM_A and RM_B on the two paths which have a transverse separation l . The shaded area is meant to represent the coronal plasma, with the gray scale conveying the strength and sign of the current density. Black regions indicate regions of large positive current density, white areas are regions of large negative current density, and gray areas are regions of zero current density. This picture is taken from numerical studies of current sheet development in MHD turbulence (Spangler 1999).

The distribution of currents shown in Figure 1 is also consistent with a picture in which the current-carrying entities are coronal loops. This case would be relevant to streamer plasmas and the closed-field part of the corona. In this case, currents flow along coronal loops, up on one footpoint and down on the other, as discussed, for example, by Chen (1989) and Chen and Schuck (2007). The lines of sight to a radio source in a differential Faraday rotation experiment could “bracket” one branch of a loop and miss the other, so the cartoon in Figure 1 reasonably represents this case as well.

The differential Faraday Rotation is defined as $\Delta RM = RM_A - RM_B$. As is illustrated in Figure 1(b), ΔRM is equivalent to the sum $RM_1 + RM_3$, where RM_1 is the same as RM_A , and RM_3 is the same as RM_B , but with the direction in which the integration is taken being reversed, i.e. from the telescope to the source. The value of ΔRM differs insignificantly from the sum of four terms, $RM_1 + RM_2 + RM_3 + RM_4$, as shown in Figure 1(c). The

reason for this is that the contributions of terms RM_2 and RM_4 are small compared to that of $RM_1 + RM_3$. There are two reasons for this. First, for the circumstances of a real coronal observation, the length of the segments 2 and 4 is small compared to 1 and 3. Second, these segments may be considered to be located far from the point of closest approach to the Sun (taken to be the middle of the diagram in all three panels), where both the plasma density and magnetic field strength are much smaller than their values near the point of closest approach.

The net result then, is that the differential Faraday Rotation is given by

$$\Delta RM = RM_1 + RM_2 + RM_2 + RM_3 = C \oint n_e \vec{B} \cdot d\vec{s} \quad (2)$$

where the last expression represents a path integral around the closed Amperian Loop defined by the segments 1,2,3, and 4 in Figure 1. The constant C is defined as $C = \frac{e^3}{8\pi^2 c^3 \epsilon_0 m_e^2} = 2.631 \times 10^{-13}$ in SI units.

Equation (2) strongly recalls Ampere’s Law, but the obvious difference is that in the present case the integrand in (2) is not just the magnetic field, but the product of the plasma density and the magnetic field. If we make the assumption that the measured ΔRM is dominated by a region in which the plasma density is relatively uniform, and given by a value $\bar{n}$, we then have

$$\Delta RM = C \oint n_e \vec{B} \cdot d\vec{s} \simeq C \bar{n} \oint \vec{B} \cdot d\vec{s} \quad (3)$$

and Ampere’s Law can be utilized. The remainder of this paper will assume the convenience of equation (3). Although we think that equation (3) is a reasonable first-order approximation to employ, it is legitimate to ask for an estimate of the associated error, at least in some plausible test cases. This matter is discussed in Section 4 below.

Given the approximation of equation (3), we can now use Ampere’s Law,

$$\oint \vec{B} \cdot d\vec{s} = \mu_0 I \quad (4)$$

where μ_0 is the permeability of free space and I is the current contained within the Amperian Loop. Use of equation (4) in equation (3) allows us to write an equation giving the coronal current in terms of the measured ΔRM ,

$$I = \frac{\Delta RM}{C \mu_0 \bar{n}} \quad (5)$$

Use of equation (5) requires an estimate of $\bar{n}$. Fortunately, there are a number of empirical expressions for the coronal plasma density as a function of heliocentric distance.

For the purposes of this paper, we utilize the following expression (Spangler 2005), which has been employed in previous analyses of Faraday Rotation observations, and is in good agreement with independent estimates.

$$n(r) = 1.83 \times 10^{12} \left(\frac{r}{R_{\odot}} \right)^{-2.36} \text{ m}^{-3} \quad (6)$$

For purposes of simplifying the subsequent formulas, the index 2.36 will be rounded off to 2.5.

The density given by equation (6) would be the maximum that would be measured along a line of sight with a dimensionless impact parameter $R_0 = \frac{r_{min}}{R_{\odot}}$, where r_{min} is the smallest heliocentric distance along the line of sight. The density $\bar{n}$ in equation (5) represents an average over the Amperian Loop defined by the observations. Therefore, for a given impact parameter R_0 , $\bar{n}$ should be less than $n(R_0)$ by some factor which is defined by $\bar{n} = \alpha n(R_0)$. With these parameterizations, substitution of (6) into (5) yields the formula that will be used in Section 3,

$$I_{obs} = 1.65 \times 10^6 \frac{(\Delta RM) R_0^{2.5}}{\alpha} \text{ Amperes} \quad (7)$$

The subscript “obs” on the current indicates that it is an estimate computed from observable quantities. Before leaving this section, it is worthwhile to emphasize the fundamental fact that the current given by equation (7) is the sum of all currents, positive and negative, within the Amperian Loop. It is thus obviously possible for strong currents to be flowing, but the total current given by equation (5) or (7) to be zero.

The approach described above requires that the region which makes the dominant contribution to the Faraday rotation measure be sufficiently uniform to allow definition of a meaningful mean density $\bar{n}$. As will be seen in Section 4.2 below, modest variations in plasma density do not result in a significant quantitative error when equations (5) or (7) are applied. However, larger errors could occur if one line of sight passes through a physically distinct region relative to the other. Whether this situation is pertinent will be determined by future observations which apply the technique described in this paper, and benefit from independent diagnostics of the plasma along the line of sight.

3. Observational Implementation and Estimates of Coronal Currents

We use data from the VLA observations of 3C228 in August, 2003. Further details of the observations and data analysis are given in Spangler (2005) and Spangler et al (2007). The characteristics of the observations relevant to the present discussion are given at the beginning of Section 2 of the present paper.

Figure 3 of Spangler (2005) shows the Faraday Rotation ² time series for 3C228 on August 16, 2003. The main feature of those observations was a large change in rotation measure over the 8 hour duration of the observing session. The data shown there distinguished measurements of the Faraday rotation to three source components, the two hot spots and a portion of the radio jet which lies between the hot spots (see Figure 2 of Spangler 2005). Similar data are available for observations of 3C228 on August 18 (Spangler et al 2007).

These data have been used to calculate time series of ΔRM for both observing sessions. These data are shown in Figure 2, and are the differences in the rotation measures to the two hot spots of 3C228. The values of ΔRM plotted have used the data at both frequencies of observation, 1465 and 1665 MHz. In the case of the observations of August 18, 2003, only data from the last 4 hours of the observing session are used. Prior to this time, elevated system temperatures due to the proximity of the Sun caused a substantial loss in data quality. The most prominent signal in the data for either day is the “event” at 19h UT on August 16. This was discussed in Spangler (2005) and evidence for its credibility is given in Spangler et al (2007). The differential Faraday Rotation had a maximum absolute value of 6.52 ± 0.77 rad/m² in the scan at 19h UT, and declined over the next two scans to a value around zero.

The observations for August 18, when the line of sight passed closer to the Sun, show no such prominent event. The final scan shows a marginally significant detection of $\Delta RM = 1.15 \pm 0.42$ rad/m². Other measurements on this day are consistent with zero.

Returning to the data for August 16, the three hours prior to the 19h event show measurements consistent with $\Delta RM = 0 \pm 2$ radian/m². These data can be used to set a significant upper limit to the current within the lines of sight. Data after 20h UT on August 16 appear to show significant ΔRM values, although with no indication of a systematic mean value. We do not believe these data show good evidence for differential Faraday Rotation during this interval, because the errors are calculated from a propagation of known radiometer noise errors, and may (and probably do) underestimate the true errors due to solar interference.

The final observational point to be made is that the ΔRM event at 19h on August 16 occurred at a time when the rate of change of the rotation measure increased (see Figure 3 of Spangler 2005). As mentioned in that paper, the line of sight during this time was moving deeper into a coronal streamer, so this feature may not be typical of coronal properties.

²The rotation in the polarization position angle, which is the quantity measured. It is given by the product of the rotation measure and the square of the observing wavelength.

In particular, it may reveal the system of currents associated with a streamer rather than a network of turbulent current sheets as illustrated in Figure 1, and which constitutes the model in the calculations to be briefly described in Section 5.

The above data can be used in equation (7) to obtain estimates for the electrical current. The results are given in Table 1, in which results are given for 3 periods during August 16 and 18, 2003. In the calculations used here, we have assumed a value of α (defined above, just prior to equation (7)) of $\frac{1}{2}$. In Table 2 we use the absolute value of ΔRM , since its sign, and the corresponding sign of I , are not considered in this paper.




Fig. 1.— Illustration of the measurement of coronal currents. Lines of sight through the corona are represented by dotted lines, with the arrow indicating the direction in which the integral is taken. The square shaded area represents the coronal plasma, with the gray scale indicating the coronal current density. Positive current density is black, and negative is white. (a) Illustrates the measurements of the Faraday rotation measure along two closely-spaced lines of sight, yielding RM_A and RM_B . (b) Illustrates the differential rotation measure $\Delta RM \equiv RM_A - RM_B$. It is equivalent to the sum of RM_1 and RM_3 . (c) The differential Faraday rotation measurement is very nearly the same as the sum of the four parts $RM_1 + RM_2 + RM_3 + RM_4$, since the “end pieces” RM_2 and RM_4 contribute negligibly to the sum. The near-equality of the sum of the four segments and the differential Faraday Rotation measurement means that ΔRM is essentially equal to the path integral of $n\vec{B}$ around the Amperian Loop shown in panel (c). The gray scale representation of the coronal plasma is taken from Figure 6 of Spangler (1999).




Fig. 2.— Differential Faraday Rotation data for August 16, 2003 (left panel) and August 18, 2003 (right panel). The measurements show the difference in the rotation measures to the hot spots of the radio source 3C228, which are separated by 46 arcseconds (33,000 km in the corona). The smaller number of measurements on August 18 results from reduced data quality prior to 20h UT, due to solar interference.

From Table 1 it can be seen that detectable differential Faraday rotation requires currents of $10^8 - 10^9$ Amperes. Upper limits deduced from present radioastronomical observations will then be in this range as well.

Before leaving this section, we again emphasize that differential rotation measures ΔRM of even a few radians/m² are rare in our two August, 2003 observing sessions spanning a total observing time of 16 hours. The apparent detection of coronal currents on August 16 was an isolated event, and may be associated with a coronal streamer. The event on August 18 is of marginal statistical significance. Consequently, the typical magnitude of differential Faraday rotation through the corona at the heliocentric distances characteristic of these observations is probably less than that of the event on August 16, with net currents which are also correspondingly smaller.

4. Validity of the Constant Density Approximation

In this section we discuss the validity of the approximation employed in equation (3). The motivation in adopting it is that it is a reasonable, first order approximation which permits application of Faraday’s Law. The discussion in this section is organized into four topics.

4.1. The Nature of Coronal Current Sheets

It is obvious that until we have a better understanding of the physical nature of coronal current sheets, including their compressibility and structure, we will be unable to determine how close (3) is to an equality. The most obvious counterexample to (3), that in which the plasma density varies in a constant magnetic field, is not a satisfactory model for coronal plasma structures. In general, a spatial gradient in the plasma density results in a similarly directed gradient in the pressure. If the plasma structures are in approximate mechanical

Table 1. Results on Coronal Currents

Date	UT Time (hours)	ΔRM (rad/m ²)	I (Amperes)
Aug. 16	15.5 - 18.5	≤ 2.0	$\leq 7.7 \times 10^8$
Aug. 16	19.0	6.52 ± 0.77	$(2.50 \pm 0.30) \times 10^9$
Aug. 18	22.7	1.15 ± 0.42	$(2.34 \pm 0.86) \times 10^8$

equilibrium, this pressure gradient must be balanced, and the most obvious possibility is a $\vec{J} \times \vec{B}$ force, where $\vec{J}$ is the current density. Thus currents will be necessary where there are plasma density variations, and while the two sides of equation (3) might not be equal, they should be of comparable magnitude.

A case in which the above considerations are not valid would be if differential Faraday rotation signals are due to parallel-propagating slow magnetosonic waves, for which there are density variations but no field variations. There is also the possibility that density variations are associated with anticorrelated temperature variations, so that pressure variations are absent, and the need for currents to achieve mechanical equilibrium is removed.

4.2. The Z-Pinch as a Model for Coronal Current Filaments

In this subsection we adopt a simplified model for coronal current filaments to explicitly calculate the error associated with the approximation (3). The model we adopt is the Z-Pinch (Krall and Trivelpiece 1973; Gurnett and Bhattacharjee 2005) which is a solution to the equation for mechanical equilibrium

$$\vec{J} \times \vec{B} - \nabla p = 0 \quad (8)$$

in cylindrical symmetry, where p is the plasma pressure. In the Z-Pinch, current can flow in the z direction (coordinate perpendicular to the plane of the page in Figure 1, not the coordinate of the observational line of sight defined in eq.[1]) and θ direction (azimuthal direction in the plane of Figure 1), and thus balance pressure gradients in the radial direction in this plane. The Z-pinch thus resembles the current sheets shown in Figure 1, with the exception that we now approximate these current filaments as having circular symmetry.

Although the Z-Pinch is an unstable equilibrium (Krall and Trivelpiece 1973; Gurnett and Bhattacharjee 2005), it is a well-studied model for current-carrying plasma structures, and one can imagine compressible astrophysical turbulence as consisting of such structures forming, persisting for a while, then dispersing with subsequent reformation.

We adopt the common simplification of ignoring the θ component of the current, which is equivalent to assuming no radial gradients in the z component of the magnetic field. Equation (8) then yields the following differential equation for the relation between the plasma pressure and the θ component of the magnetic field

$$\frac{B_\theta}{r} \frac{d(rB_\theta)}{dr} = -\mu_0 \frac{dp}{dr} \quad (9)$$

where r in this section is the radial coordinate in a cylindrical coordinate system, not

the heliocentric distance as used in Section 2. It is B_θ that causes the differential Faraday rotation in this model. In discussions of the Z-Pinch (Krall and Trivelpiece 1973; Gurnett and Bhattacharjee 2005), one prescribes one of the fields p or B_θ , then uses equation (9) to determine the other. This procedure is well-suited to our present task, which is to see how spatial gradients in the plasma density of a plausible plasma structure affect differential Faraday rotation.

We assume an isothermal approximation for the plasma equation of state, which will give a larger density change for a given pressure change than adiabatic equations of state.

$$p(r) = c_s^2 \rho = c_s^2 m_H n(r) \quad (10)$$

where c_s is the sound speed, ρ is the mass density, m_H is the mass of a hydrogen atom, and $n(r)$ is the number density of atoms. We assume a pure hydrogen plasma. Substituting equation (10) into (9), and making the change of variable $B_\theta \rightarrow rB_\theta \equiv f$, equation (9) becomes

$$\frac{df^2(r)}{dr} = -2\gamma r^2 \frac{dn(r)}{dr} \quad (11)$$

with $\gamma \equiv \mu_0 c_s^2 m_H$, which is directly integrable. The solution is

$$f^2(r) = (rB_\theta(r))^2 = -2\gamma \int_0^r r^2 \frac{dn}{dr} dr \quad (12)$$

We adopt a model for the density profile which is enhanced in the current channel,

$$n(r) = \bar{n} + n_1 e^{-(r/a)} \quad (13)$$

with a being the specified radial extent of the current filament. We substitute equation (13) into (12), and change variable of integration from $r \rightarrow \xi \equiv \frac{r}{a}$. The solution then becomes

$$f^2(r) = (rB_\theta(r))^2 = 2\gamma n_1 a^2 F(\xi) \quad (14)$$

$$F(\xi) \equiv \int_0^\xi \xi^2 e^{-\xi} d\xi = 2 - (\xi^2 + 2\xi + 2)e^{-\xi} \quad (15)$$

$$B_\theta(r) = \frac{\sqrt{2\mu_0}}{r} \sqrt{\Delta p a} \sqrt{F(\xi)} \quad (16)$$

where $\Delta p \equiv c_s^2 m_H n_1$. Equation (16) also yields the expression for the current density

$$J_z(r) = \left(\frac{1}{a}\right) \sqrt{\frac{\Delta p}{2\mu_0}} G(\xi) \quad (17)$$

$$G(\xi) \equiv \frac{\xi e^{-\xi}}{\sqrt{F(\xi)}} \quad (18)$$

We now express the different fields in the current filament in terms of common parameters which characterize the host plasma and the current channel. The plasma pressure is, from above,

$$p = c_s^2 m_H (\bar{n} + n_1 e^{-\xi}) \quad (19)$$

We define the “modulation index” $\epsilon \equiv \frac{n_1}{\bar{n}} = \frac{\Delta p}{p_0}$, so that

$$n = \bar{n}(1 + \epsilon e^{-\xi}) \quad (20)$$

$$p = p_0(1 + \epsilon e^{-\xi}) \quad (21)$$

With $p_0 \equiv c_s^2 m_H \bar{n}$. We now relate the plasma pressure to the magnetic pressure via

$$p_0 = \beta \left(\frac{B_0^2}{2\mu_0} \right) \quad (22)$$

$$\Delta p = \epsilon \beta \left(\frac{B_0^2}{2\mu_0} \right) \quad (23)$$

Where β is the conventional plasma β , here defined as the ratio of gas pressure to magnetic pressure, and B_0 is the mean magnetic field. Using equation (23) in the equation for B_θ gives

$$B_\theta = B_0 \sqrt{\epsilon \beta F_2(\xi)} \quad (24)$$

$$F_2(\xi) \equiv \frac{\sqrt{F(\xi)}}{\xi} \quad (25)$$

With these expressions, we can now calculate both sides of the near-equality (3). The integral is taken in a clockwise sense, as indicated in Figure 1(b), on the Amperian Loop shown in Figure 1, with the convention that the origin of the polar coordinate system is midway between the integration paths A and B. The results of the calculation below are not qualitatively changed if the center of the current filament ($r = 0$) is not midway between the two linear paths of integration A and B. The segments of the Amperian Loop connecting paths A and B (segments 2 and 4 in Figure 1(c)) are assumed to lie at infinity and are neglected in the calculation below.

The integral around the Amperian Loop is then given by symmetry considerations as

$$\oint n \vec{B} \cdot d\vec{s} = 4 \int_b n \vec{B} \cdot d\vec{s} \quad (26)$$

where the path segment b corresponds to $\frac{\pi}{2} > \theta > 0$. In what follows we evaluate the integral over path b. Given the expression for B_θ , etc developed above, the integrand of equation (3) is

$$n \vec{B} \cdot d\vec{s} = n(\xi) B_\theta(\xi) \hat{e}_\theta \cdot \hat{e}_x dx \quad (27)$$

where $\hat{e}_x$ is a unit vector in the the x direction (horizontal axis in Figure 1), and $\hat{e}_\theta$ is the unit vector in the direction of increasing θ . The dot product of the unit vectors is $\hat{e}_\theta \cdot \hat{e}_x = -\cos \bar{\theta}$, where $\bar{\theta}$ is the complement of θ .

An important variable is the perpendicular distance R of the integration path from the center of the current channel. In terms of our observations, $2R$ would be the separation l between the lines of sight through the corona. The incremental path element $dx = R \sec^2 \bar{\theta} d\bar{\theta}$, and the variable ξ is $\xi = \frac{R}{a} \sec \bar{\theta}$. Substituting all of these relations into (27) and forming the integral, we have

$$\oint n \vec{B} \cdot \vec{ds} = 4 \int_0^{\pi/2} n(\xi) B_\theta(\xi) R \sec \bar{\theta} d\bar{\theta} \quad (28)$$

Substituting equations (20) and (24) into (28), we have

$$\oint n \vec{B} \cdot \vec{ds} = 4n_0 B_0 \sqrt{\epsilon \beta} R \left[\int_0^{\pi/2} F_2(\xi) \sec \bar{\theta} d\bar{\theta} + \epsilon \int_0^{\pi/2} e^{-\xi} F_2(\xi) \sec \bar{\theta} d\bar{\theta} \right] \quad (29)$$

$$= 4n_0 B_0 \sqrt{\epsilon \beta} R [S_a + \epsilon S_b] \quad (30)$$

$$= 4n_0 B_0 \sqrt{\epsilon \beta} R S_a \left[1 + \epsilon \frac{S_b}{S_a} \right] \quad (31)$$

Equations (29) and (30) define the integrals $S_a \left(\frac{R}{a} \right)$ and $S_b \left(\frac{R}{a} \right)$. An appealing feature of equation (31) is that the leading multiplicative term on the right hand side is the value of $\oint n \vec{B} \cdot \vec{ds}$ for the case of uniform density $\bar{n}$. The second term in brackets in (31) is therefore the correction due to the change in density in the Z-Pinch. It is directly proportional to the density modulation index ϵ .

The parameter $\Gamma \equiv \frac{\epsilon S_b}{S_a}$ gives the fractional error in approximating $\oint n \vec{B} \cdot \vec{ds}$ by $\bar{n} \oint \vec{B} \cdot \vec{ds}$. Obviously, this is equal to the fractional error in the current inferred from the radioastronomical technique which is the subject of this paper. Given the definition of the density modulation index ϵ , one would expect it to be less than unity, say $\epsilon \leq 0.50$, although this is not strictly required in the definition of equation (20). In any case, it is a parameter which might be observationally accessible. Given an adopted value of ϵ , the correction factor which relates the true current in a plasma to that retrieved by the technique of Section 2 is then dependent on the ratio $\frac{S_b}{S_a}$. The integrals S_a and S_b were evaluated numerically with a *Mathematica* notebook. The results for a range of values of $\frac{R}{a}$ are given in Table 2.

The conclusions to be drawn from the calculations summarized in Table 2 are clear. Even for lines of sight which pass deep in the interior of the current channel, the ratio $\frac{S_b}{S_a} < 1$ by a substantial factor. The fractional error $\Gamma = \frac{\epsilon S_b}{S_a}$ will be smaller still, even for relatively large compression factors of $\epsilon \simeq 0.5$. For example, if the line of sight passes within $R = 0.1 - 0.2a$ of the center of the filament, and the compression factor $\epsilon = 0.50$, the error

in the current retrieved by the method of Sections 2 and 3 would be only 17 - 19 %. This is completely negligible in the context of this paper, and indicates that the method described here would be quite accurate.

For lines of sight that pass further from the center of the filament, the fractional error becomes smaller still, approaching zero as $\frac{R}{a} \gg 1$, as indeed it must.

4.3. Analysis of Measurements from the MST

In this subsection, we consider a test of our method with data from the Faraday rotation experiment on the Madison Symmetric Torus (MST) plasma machine (Brower et al 2002; Ding et al 2003). This approach has the advantage of using data from a real plasma, in contrast to the mathematical ideal of Section 4.2. In addition, as a laboratory experiment, it is diagnosed in a way that the solar corona cannot be.

In some sense, use of the MST data assures corroboration of the technique. The goal of the experiments was to measure currents flowing in the MST, and those experiments motivated the current investigation. However, the analysis of Brower et al (2002) and Ding et al (2003) utilized more experimental information than is available to us, and (at least in part) applied more sophisticated methods of processing the data, which benefitted from a-priori information on the structure of the toroidal plasma.

In this subsection, we apply the same formulas used in Sections 2 and 3 for 3C228 to published data from the MST (Brower et al 2002), and compare the retrieved current with that known to flow in the MST device. Brower et al (2002) report that a current of 400 kA flows in the MST during their experiments, and that the mean density (which we adopt for $\bar{n}$) is 10^{19} m^{-3} . The wavelength of the polarized laser beam used to probe the plasma was 432 nm.

We utilize the data shown in Figure 1 of Brower et al (2002), which shows Faraday rotation of about 5° for the chord at an offset distance of -17 cm (corresponding to the parameter R defined in the previous subsection), and Faraday rotation of about $-3^\circ.5$ for the chord at $R = +21$ cm. We thus have differential Faraday rotation of $\Delta\chi \simeq 8^\circ.5$, corresponding to a differential rotation measure $\Delta RM \simeq 8.0 \times 10^5 \text{ rad/m}^2$ between two lines of sight separated by 0.38 meters. Substitution of this value for ΔRM into equation (5) yields a current of 241 kA.

This value itself, resulting from the most straightforward application of the methods used in the study of the corona, is in reasonably good agreement with the known value.

It is within a factor of two of the known, measured current in the MST device. Similar precision in the measurement of coronal currents would be cause for delirium. However, we can improve agreement between remotely inferred currents in the MST, using equation (5) of this paper, and the known value for the current.

The laser polarimeter system of Brower et al (2002) and Ding et al (2003) utilized a number of chords with different values of R , so the current profile can be retrieved. In fact, the change in the functional form of the current density during a sawtooth crash on MST was a main result of Brower et al (2002). The inferred toroidal current density profile $J_z(r)$ is plotted in Figure 4 of Brower et al (2002), and shows that the current channel has a greater radial extent than 0.2m. Accordingly, the estimate of I immediately above must be an underestimate. To approximately calculate the current within the two outer chords displayed in Figure 1 of Brower et al (2002), we represent the current density profile $J_z(r)$ by a Gaussian,

$$J_z(r) = J(0)e^{-(r/a)^2} \quad (32)$$

We use plot 4(b) of Brower et al (2002) and the profile measured before a sawtooth crash, for which we estimate values of $J(0) = 1.93 \times 10^6$ A/m², and $a = 0.29$ m. The total current within two lines of sight separated by a distance $2R$ with the center of the current filament midway between the lines of sight is

$$I(R) = \pi J(0) a^2 \text{Erf}(X) \quad (33)$$

where $\text{Erf}(X)$ is the Error Function of $X \equiv \frac{R}{a}$. Using the above values of $J(0)$ and a , and with $R = 0.19$ m, equation (33) gives $I(R = 0.19\text{m}) = 331\text{kA}$. This value is, as expected, in better agreement with the value inferred from equation (5), and indicates a 37 % error between the “remote sensing” and “true” value, a difference which includes possible systematic errors in the plasma physics experiment, which are obviously beyond the scope of this paper. In any case, the degree of agreement supports the credibility of the technique discussed in this paper.

The work of Brower and Ding demonstrates that future coronal differential Faraday rotation observations, with measurable ΔRM on several lines of sight to a background radio source, could measure not only the total current but the profile of coronal current density, as is done in the MST. To summarize the results of this and the preceding subsection, investigation of a simplified analytical model of coronal current filaments and experimental results from a laboratory plasma have suggested that the approximation of equation (3) results in errors in the retrieved current of, at most, tens of percent. Such an error does not detract from the significance of our results for coronal physics.

4.4. Prospects for Improved Modeling of Coronal Currents

The analysis of Section 4.2 would be improved through the use of realistic models for the compressive current systems which exist in the solar corona. As in the case of Sections 4.2 and 4.3, the goal will be to determine the error associated with assuming a constant density in the plasma, so that the associated error in the retrieved current can be estimated.

A promising set of existing calculations is comprised of the recent coronal simulations of Gudiksen and Nordlund (2005) and Peter et al (2006), which are specifically directed to studies of Joule heating of the corona. The calculations presented in those papers are three dimensional and give the vector magnetic field, plasma density, and current density at all locations in the simulation. The Faraday rotation measure along different lines of sight through the simulation could be calculated and the relationship between differential Faraday rotation and current properties investigated.

The investigations to date by Gudiksen, Peter, and collaborators have concentrated on the corona at much smaller heliocentric distances than probed by our VLA observations. However, future calculations could presumably be carried out to simulate conditions at heliocentric distances of $5 - 10R_{\odot}$. Furthermore, even the present calculations which have been described in the literature could be used to examine the general relation between differential Faraday rotation and current filaments in a simulated environment patterned after the corona.

5. Implications for Coronal Heating

Having provided observational estimates of coronal currents in Section 3 of this paper, there arises the obvious question as to whether these currents can provide the requisite heating of the corona via Joule heating. Unfortunately, it is not straightforward to answer this question. The volumetric heating rate, which must be the basis of such a discussion, is ηJ^2 where η is the electrical resistivity of the plasma, and J is the current density. To obtain the average coronal heating rate in the region probed by our observations, one must estimate the geometry, extent, and number of the current sheets in a given volume of the corona, in order to estimate both J from a measurement of I , and also to calculate the filling factor of the volume in which heating is going on. This obviously requires a model for the current sheets. Furthermore, and most importantly, an expression for the resistivity η must be chosen.

We carried out a calculation for the average volumetric heating rate, using two models for the nature of the current sheets responsible for the differential Faraday rotation. These

calculations envisioned the corona as illustrated in Figure 1, with thin, long (along the large-scale coronal field) sheets, in which the sign of the current density would vary from one sheet to the next. One of the two models assumed that there was equal probability of positive and negative current density, as would be expected in a true turbulent situation, and in which the measured differential Faraday rotation is due to a statistical fluctuation in the number of positive and negative sheets within the Amperian Loop. The second model assumed a statistical preference for one sign of the current density.

Formulas were derived for effective volumetric heating rates in both cases. These heating rates depend on assumed properties of the current sheets, such as thickness, lateral extent, extension along the large scale field, etc. The heating rates also depend crucially on the resistivity. Plausible estimates could be obtained for all the geometric properties of these current sheets. For the resistivity, we used the Spitzer resistivity (Gurnett and Bhattacharjee 2005) which is based on fully-understood principles and may be assumed to be a lower limit to the true coronal resistivity. The volumetric heating rate for either model was roughly 6 orders of magnitude less than that estimated by Cranmer and Ballegoijen (2005) as necessary to explain heating in the relevant part of the corona. This calculation indicates that either the currents which are reported here are irrelevant for coronal heating, or that they do play a role in coronal heating, but that the effective resistivity of the coronal plasma at heliocentric distances of $5 - 10R_{\odot}$ is approximately 6 orders of magnitude greater than the Spitzer value. A document with a detailed description of these calculations is available from the author.

6. Summary and Conclusions

1. We have pointed out that a type of polarization measurement (differential Faraday Rotation measurements on an extended radio source) which can be done with the Very Large Array can yield estimates of electrical currents in the solar corona. The technique could, in principle, be done with some other radio telescopes as well. This technique is an astronomical adaptation of a diagnostic used in fusion plasmas, and described by Brower et al (2002) and Ding et al (2003).
2. The technique has been applied to observations of the radio source 3C228 which were made on two days in August 2003, when the radio source was viewed through the corona at “impact parameters” of 6.7 and $5.2R_{\odot}$. Detectable differential Faraday rotation was detected in relatively brief “events” on both days, yielding estimated currents of 2.5×10^9 and 2.3×10^8 Amperes, respectively. Another interval of high quality data on one of the days yielded an upper limit to the differential Faraday rotation, and a corresponding upper limit to the current of 8×10^8 Amperes.

3. The technique as defined in this paper is dependent on an approximation in which the plasma density is assumed constant in the region within the Amperian Loop which makes the dominant contribution to the rotation measure. The validity of this approximation was investigated in two ways in Section 4. First, the value of $\oint n \vec{B} \cdot d\vec{s}$ was compared with $\bar{n} \oint \vec{B} \cdot d\vec{s}$ for a Z-Pinch model in which the plasma density is modified in the current filament. Second, the formula used in our radioastronomical observations (eq.[5]) was applied to measurements made with an infrared laser polarimeter on the MST plasma machine (Brower et al 2002; Ding et al 2003), and used to estimate the current, which could also be measured independently. The values agreed to 37 %.
4. We explored the implications of the measured currents for Joule heating of the corona. The magnitude of the estimated, average volumetric heating rate is rendered uncertain by poorly-known properties of the current sheets or filaments, and by the value of the coronal resistivity. Use of the Spitzer resistivity leads to a value for the volumetric heating rate approximately 6 orders of magnitude less than the value indicated by observations. We conclude that either these currents are irrelevant for coronal heating, or that the true resistivity in the corona exceeds the Spitzer value by several orders of magnitude. Resolution of this matter obviously lies in a better understanding of the resistivity in a collisionless plasma.

This work was supported at the University of Iowa by grants ATM03-11825, and ATM03-54782 from the National Science Foundation. I thank W.X. Ding and D.L. Brower of the University of California, Los Angeles, for interesting and helpful discussion about Faraday rotation diagnostics of the current profile on the MST Reversed Field Pinch device of the University of Wisconsin.

REFERENCES

- Bird, M.K. and Edenhofer, P. 1990, in Physics of the Inner Heliosphere II, R. Schwenn and E. Marsch, ed., (Springer-Verlag:Berlin), p13
- Brower, D.L. et al. 2002, Phys. Rev. Lett. 88, 185005
- Chen, J. 1989, ApJ 338, 453
- Chen, J. and Schuck, J.W. 2007, Sol. Phys.(submitted)
- Cranmer, S.R. and Ballegooijen, A.A. 2005, ApJS 156, 265

- Ding, W.X. et al. 2003, Phys. Rev. Lett. 90, 035002
- Gudiksen, B.V. and Nordlund, Å. 2005, ApJ 618, 1020; *Erratum*: ApJ 623,600
- Gurnett, D.A. and Bhattacharjee, A. 2005, Introduction to Plasma Physics, Cambridge University Press
- Hollweg, J.V., Bird, M.K., Volland, H., Edenhofer, P., Stelzried, C.T., and Seidel, B.L. 1982, J. Geophys. Res. 87, 1
- Ingleby, L.D., Spangler, S.R., and Whiting, C.A. 2007, ApJ(in press, Oct. 10, 2007; astro-ph/0701538)
- Krall, N.A. and Trivelpiece, A.W. 1973, Principles of Plasma Physics, McGraw-Hill
- Kraus, J.D. 1966, Radio Astronomy, McGraw-Hill, p144
- Mancuso, S. and Spangler, S.R. 1999, ApJ 525, 195
- Mancuso, S. and Spangler, S.R. 2000, ApJ 539,480
- Parker, E.N. 1972, ApJ 174,499
- Peter, H., Gudiksen, B.V., and Nordlund, Å. 2006, ApJ 638, 773
- Prager, S.C. 1999, Plasma Phys. Control. Fusion 41, A129
- Sakurai, T. and Spangler, S.R. 1994, ApJ 434, 773
- Spangler, S.R. 1999, ApJ 522,879
- Spangler, S.R. 2005, Space Sci. Rev. 121,189
- Spangler, S.R., Spitler, L.G., Miralles, M.P., Cranmer, S.R., Raymond, J.C., and Kasper, J.C. 2007, (in preparation)

Table 2. Validity of Constant Density Approximation

$\frac{R}{a}$	S_a	S_b	$\frac{S_b}{S_a}$
0.1	1.682	0.645	0.383
0.2	1.585	0.542	0.342
0.4	1.443	0.397	0.275
0.6	1.331	0.297	0.223
0.8	1.237	0.224	0.181
1.0	1.154	0.169	0.147
1.4	1.015	0.098	0.096
1.8	0.902	0.057	0.063

February 5, 2008

Dr. Timothy Bastian, Editor
The Astrophysical Journal
National Radio Astronomy Observatory
520 Edgemont Road
Charlottesville, Virginia 22903

Dear Tim,

I have just submitted to the Astrophysical Journal server the *revised version* of my paper, “A Technique for Measuring Electrical Currents in the Solar Corona”, manuscript # 71459. To review the history of this paper, it was submitted to the Astrophysical Journal on February 16 of this year. I received the report of the referee (which included comments both by the referee and an observational colleague) on February 26. Your assessment of the referee’s report was that the paper was “unacceptable in its current form”.

I was encouraged to revise and resubmit this paper for two reasons. First, the referee was favorably inclined to the basic idea of the paper, saying that “the idea is explained clearly, and the paper should be published in some form”, and “the central idea of this paper should be published”. Second, it seemed clear to me that the referees’ objections were mainly in two areas which could be addressed. It is my hope that I have done so.

The revised paper is a major revision, and involves a substantial amount of new work, undertaken specifically to address the criticisms of the referees. To guide the referees in assessing the revised version, and as an overview to what I have done, the first three sections (up to and including the calculation of currents from the observations) are only slightly revised from the original, in ways which incorporate suggestions from the referees. A completely new Section 4 has been inserted to address the validity of the constant density approximation. Finally, the revised Section 5 is, as recommended by both referees, a “severely abbreviated” version of the calculation of the heating rate by Joule heating of these currents.

The criticisms of the primary referee are contained in the “Three Disappointments”, a term that suggests a dark variation on an entree in a Chinese restaurant. I hope my responses to the “Three Disappointments” fully and positively respond to these suggestions. There are also the criticisms and suggestions by the observational colleague. The observational colleague’s remarks consist of a preamble, three enumerated comments on the method, and two comments on the interpretation.

In what follows, I give my response to all of these points.

The “Three Disappointments”

(1) The referee wishes to have a discussion of the approximation/assumption of pulling the density outside the path integral in equation (3) (both versions). Although I thought this was a reasonable first order approximation, the referee obviously has a legitimate issue here. In response to this point, I have included an entire new section, the present Section 4 of the paper, entitled “Validity of the Constant Density Approximation”. There are four points brought up in this section. (A) Until we have a better idea of the structure and thermodynamics of coronal irregularities, we cannot really answer this question. However, I do make the point that if there are density fluctuations, there will generally (but not in all cases) be corresponding electrical currents. (B) The main part of this section, Section 4.2, is a quantitative discussion of this approximation for a model of coronal current filaments based on the Z-Pinch. The Z-Pinch is a current-carrying filament which confines an enhanced density by $\vec{J} \times \vec{B}$ forces. This model is analysed quantitatively and a parameter is obtained (the parameter Γ introduced in the last paragraph on p15) which gives the error due to the approximation involved in equation (3) of this paper. The quantitative results, given in Table 2 of the revised version, show that the error should be of order 20 % in a fairly unfavorable case, and much less otherwise. Obviously, we have no assurance that the coronal current sheets are structured like Z-Pinches, but the point here is that in the case of a plausible facsimile for these loops, the error in the constant density approximation is very small. (C) I took data from the figures in the MST Reversed Field Pinch experiments (the experiments which inspired the development of this technique) and applied my formulas which were used for the observations of radio galaxies through the corona. This gave a “remote sensing” estimate of the current in the machine, which is independently known. In Section 4.3 of the revised paper, I describe these calculations, and show that the “retrieved” current is within 37 % of the known current in the plasma device. So this technique is shown to work (at the level of a few tens of percent) in the case of a “real” plasma. (D) I point out that this issue could be settled definitively by calculating the differential Faraday rotation along paths through numerical simulations of the corona in which Joule heating is playing a major role, and I reference the papers of Gudiksen, Nordlund, and Peter. This would be a very straightforward thing to do, and I hope my paper (if it is published) could encourage such a project.

(2) The referee asked that the possibility be mentioned that the currents could be those directed along coronal loops rather than the set of turbulent current sheets that I had envisioned. In the revised paper, I have included a paragraph of new text, third complete paragraph, p4, in which I discuss this possibility and reference two papers of Jim Chen’s which specifically discuss (and illustrate in the figures) electrical currents flowing on coronal loops.

(3) The referee obviously did not like the section on estimates of Joule heating at a rate

determined by the Spitzer resistivity, and stated in his or her recommendation # 4 that “the discussion of coronal heating by the putative current should be severely abbreviated or deleted entirely”. I was very reluctant to do this. I anticipated at the outset that Joule heating at the Spitzer rate would be far smaller than real rates, but I felt it was important to actually demonstrate that this is the case. Furthermore, although the final expression for the heating rate involved many “imponderables”, I think it important to go through the exercise and identify what parameters are most important in determining the true heating rate. The referee may be interested in knowing that I undertook this calculation at the encouragement of Peter Gary.

Nonetheless, Peter Gary was not the referee of this paper, so I felt I had to modify my paper to satisfy the current referee. I have therefore chosen the “severely abbreviated” route, and the new Section 5 consists of only three paragraphs of text and no equations, and summarizes the results of the old Section 4 of the original paper. I also mention that the details of this calculation are available from the author. I really think I have to do this. In view of the prominence of Joule heating theories for coronal heating, a report which describes how some observational handle can be had on currents has to say something about how these numbers relate to Joule heating calculations.

Comments of the Observational Colleague

The preamble to this person’s remarks makes the entirely valid point that the main “event” on August 16, 2003 occurred at a time when the line of sight to the radio source was passing into a streamer, so the currents which are being detected might be rare and associated with a single structure, rather than due to turbulent current sheets of the sort I have envisioned.

I agree with everything the referee says here, and believe I clearly described the situation in the original version of the paper. However, to emphasize this point, I have inserted a paragraph of new text on p11, second complete paragraph, last paragraph of Section 3, which clearly indicates that most of the time, the differential rotation measure was less than that observed at 19h on August 16. In this case, as I indicate, the sort of analysis here could be used to set upper limits to the current. With regard to the marginal nature of the ΔRM on August 18, I stated the error on the measurement in the original version as well as the current version, so I did “put my cards on the table”. Nonetheless, to make this completely clear I have appended the modifier “marginal” when referring to it in the revised version of the paper (see, for example, the description of this measurement in the last paragraph of Section 3). Once again, one may certainly use this measurement for an upper limit to the current.

(4) The first comment in “issues with the method” notes that the differential Faraday rotation measurement is an integral over the path through the corona. I would respond that

this is completely obvious from the equations that describe the observations, beginning with equation (1). Furthermore, Ampere’s Law actually requires that the measurement be related to an integral around the whole curve (the Amperian Loop). The point that one is averaging currents that are both small and large, and of both signs does need to be kept in mind, as is pointed out in the two sentences following equation (7), revised version.

(5) The referee makes the very good point that if one line of sight goes through a region which is physically distinct from that perforated by the other LOS, a differential Faraday rotation signature would be observed which would be hard to interpret. I convey this point in a paragraph of new text inserted on p6 at the very end of Section 2. I also point out that future observations, using radio data such as mine and independent coronal diagnostics such as coronagraph observations, might indicate if this extreme inhomogeneity is affecting the observations. However, as I also point out in this new paragraph, the analysis of Section 4.2 indicates that mild density inhomogeneity does not invalidate the estimates of the current.

(6) I don’t agree with the assertion that this method is reliant on assumptions about lack of time variability. Ampere’s Law does not contain time derivatives (assuming that the displacement current is negligible), and the VLA makes observations along the two lines of sight A and B simultaneously, i.e. no time lag at all. We do not rely on Taylor’s hypothesis of frozen plasma fluctuations as is sometimes done in analysis of spacecraft data.

(7) Comments A and B, as well as their summary at the end of the report raise questions about the plausibility of parameters of the current sheets in my Joule heating calculations. The referee speculates whether the heating rate is based on a large number of contrived parameters. I can give a defense of the estimates I chose, as well as a frank “I don’t know” when it is warranted. However, since the section containing these calculations has been all but eliminated, on the basis of the recommendations of both referees, I think this point is now moot.

The new, extensive discussion of the validity of the constant density approximation, as well as the de-emphasis of the coronal heating calculation, are reflected in the revised abstract and conclusions.

This concludes my response to the referees. I hope that it is clear that I have taken their remarks seriously, and substantially modified the paper in response to their recommendations. I have done this even though I would have preferred to retain the calculation on heating. I

hope that the paper is now acceptable for publication in The Astrophysical Journal.

Sincerely yours,
Steve

Steven R. Spangler
Professor of Physics and
Astronomy